

FDA-Approved Patient Labeling

NEXPLANON® (etonogestrel implant)

Radiopaque

Subdermal Use Only

NEXPLANON® does not protect against HIV infection (the virus that causes AIDS) or other sexually transmitted diseases.

Read this Patient Information leaflet carefully before you decide if NEXPLANON is right for you. This information does not take the place of talking with your healthcare professional. If you have any questions about NEXPLANON, ask your healthcare professional.

What is NEXPLANON?

NEXPLANON is a hormone-releasing birth control implant for prevention of pregnancy for up to 5 years. The implant is a flexible plastic rod about the size of a matchstick that contains a progestin hormone called etonogestrel. It contains a small amount of barium sulfate (15 mg), so that the implant can be seen by X-ray, an ethylene vinyl acetate (EVA) copolymer (28% vinyl acetate, 43 mg) core, and magnesium stearate (0.1 mg). Your healthcare professional will insert the implant just under the skin of the inner side of your upper arm. You can use a single NEXPLANON implant for up to 5 years. NEXPLANON does not contain estrogen.



What if I need birth control for more than 5 years?

The NEXPLANON implant must be removed by the end of 5 years. Your healthcare professional can insert a new implant under your skin after taking out the old one if you choose to continue using NEXPLANON for birth control.

What if I change my mind about birth control and want to stop using NEXPLANON before 5 years?

Your healthcare professional can remove the implant at any time. You may become pregnant as early as the first week after removal of the implant. If you do not want to get pregnant after your healthcare professional removes the NEXPLANON implant, you should start another birth control method right away.


How does NEXPLANON work?

NEXPLANON prevents pregnancy in several ways. The most important way is by stopping the release of an egg from your ovary. NEXPLANON also thickens the mucus in your cervix and this change may keep sperm from reaching the egg. NEXPLANON also changes the lining of your uterus.

How well does NEXPLANON work?

When the NEXPLANON implant is placed correctly, your chance of getting pregnant is very low (less than 1 pregnancy per 100 women who use NEXPLANON for 1 year).

The following chart shows the chance of getting pregnant for women who use different methods of birth control. Each box on the chart contains a list of birth control methods that are similar in effectiveness. The most effective methods are at the top of the chart. The box on the bottom of the chart shows the chance of getting pregnant for women who do not use birth control and are trying to get pregnant.



Who should not use NEXPLANON?

Do not use NEXPLANON if you have any of the following conditions:

- Are pregnant or think you may be pregnant
- Have, or have had, blood clots, such as blood clots in your legs (deep vein thrombosis), lungs (pulmonary embolism), eyes (total or partial blindness), heart (heart attack), or brain (stroke)
- Liver disease or a liver tumor
- Unexplained vaginal bleeding
- Breast cancer or any other cancer that is sensitive to progestin (a female hormone), now or in the past
- An allergy to anything in NEXPLANON

Tell your healthcare professional if you have or have had any of the conditions listed above. Your healthcare professional can suggest a different method of birth control.

In addition, talk to your healthcare professional about using NEXPLANON if you have any of the following conditions:

- Diabetes
- High cholesterol or triglycerides
- Headaches

- Gallbladder or kidney problems
- A history of depressed mood
- High blood pressure
- An allergy to numbing medicines (anesthetics) or medicines used to clean your skin (antiseptics). These medicines will be used when the implant is placed into or removed from your arm.

Interaction with Other Medicines

Tell your healthcare professional about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using certain medicines with NEXPLANON, including those listed below, may make NEXPLANON less effective:

- aprepitant
- barbiturates
- bosentan
- carbamazepine
- felbamate
- griseofulvin
- oxcarbazepine
- phenytoin
- rifampin
- St. John's wort
- topiramate
- HIV medicines
- Hepatitis C Virus medicines

Ask your healthcare professional if you are not sure if your medicine is one listed above.

If you are taking medicines or herbal products that might make NEXPLANON less effective, you and your healthcare professional may decide to leave NEXPLANON in place; in that case, an additional non-hormonal contraceptive should be used. It is necessary to use the additional non-hormonal contraceptive for 28 days after stopping the medicine or herbal product because its effect on NEXPLANON may last that long.

When you are using NEXPLANON, tell all your healthcare professionals that you have NEXPLANON in place in your arm.

Magnetic Resonance Imaging (MRI) Safety Information

NEXPLANON is MRI safe.

How is the NEXPLANON implant placed and removed?

Your healthcare professional will place and remove the NEXPLANON implant using a minor surgical procedure in his or her office. The implant is placed just under the skin on the inner side of your non-dominant upper arm.

The timing of insertion is important. Your healthcare professional may take the following steps to guide timing of insertion:

- Perform a pregnancy test before inserting NEXPLANON
- Schedule the insertion at a specific time of your menstrual cycle (for example, within the first 5 days of your regular menstrual bleeding). If the implant is placed after the fifth day of menses, then you should use an additional contraceptive method (such as a condom) for the first 7 days after insertion.

Your healthcare professional will cover the site where NEXPLANON was placed with 2 bandages. A small bandage will be placed over the insertion site. A larger pressure bandage will be placed over the smaller bandage. Leave the larger pressure bandage on for 24 hours. Keep the smaller bandage clean, dry, and in place for 3 to 5 days.

Immediately after the NEXPLANON implant has been placed, you and your healthcare professional should check that the implant is in your arm by feeling for it.

If you cannot feel the implant immediately after insertion, the implant may not have been inserted, or it may have been inserted deeply. A deep insertion may cause problems with locating and removing the implant. When the healthcare professional has located the implant, it should be removed.

If at any time you cannot feel the NEXPLANON implant, contact your healthcare professional immediately and use a non-hormonal birth control method (such as condoms) until your healthcare professional confirms that the implant is in place. You may need special tests to check that the implant is in place or to help find the implant when it is time to take it out. If the implant cannot be found in the arm after a thorough search, your healthcare professional may use X-rays or other imaging methods on your chest.

Depending on the exact position of the implant, removal may be difficult and may require surgery.

Keep track of the date the implant is to be removed. Schedule an appointment with your healthcare professional to remove the implant on or before the removal date.

Be sure to have checkups as advised by your healthcare professional.

What are the most common side effects I can expect while using NEXPLANON?

• Changes in Menstrual Bleeding Patterns (Menstrual Periods)

The most common side effect of NEXPLANON is a change in your normal menstrual bleeding pattern. In studies, one out of ten women stopped using the implant because of an unfavorable change in their bleeding pattern. You may experience longer or shorter bleeding during your periods or have no bleeding at all. The time between periods may vary, and you may also have spotting in between periods.

Tell your healthcare professional right away if you have the following concerns:

- You think you may be pregnant
- Your menstrual bleeding is heavy and prolonged

The following additional frequent side effects also caused women to stop using the implant:

- Mood swings
- Depressed mood
- Weight gain
- Headache
- Acne

Other common side effects include the following conditions:

- Headache
- Vaginitis (inflammation of the vagina)
- Weight gain
- Acne
- Breast pain
- Viral infections such as sore throats or flu-like symptoms
- Stomach pain
- Painful periods
- Mood swings, nervousness, or depressed mood
- Back pain
- Nausea
- Dizziness
- Pain
- Pain at the site of insertion

Implants have been reported to be found in a blood vessel, including a blood vessel in the lung which can be associated with shortness of breath, cough and/or the coughing up of blood or blood-stained mucus.

These are not all the possible side effects of NEXPLANON. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

What are the possible risks of using NEXPLANON?

• Problems with Insertion and Removal

The implant may not be actually in your arm due to a failed insertion. If this happens, you may become pregnant. Immediately after insertion, and with help from your healthcare professional, you should be able to feel the implant under your skin. If you can't feel the implant, tell your healthcare professional.

Location and removal of the implant may be difficult or impossible because the implant is not where it should be. Special procedures, including surgery in the hospital, may be needed to remove the implant. If the implant is not removed, then the effects of NEXPLANON will continue for a longer time.

Implants have been found in the pulmonary artery (a blood vessel in the lung). If the implant cannot be found in the arm, your healthcare professional may use X-rays or other imaging methods on the chest. If the implant is located in the chest, surgery may be needed.

The following problems related to insertion and removal may also occur:

- Vasovagal reactions (such as a drop in blood pressure, dizziness, or fainting)
- Pain, irritation, swelling, or bruising at the insertion site
- Numbness and tingling at the insertion site
- Scarring, including a thick scar called a keloid around the insertion site
- Infection
- Scar tissue may form around the implant making it difficult to remove
- The implant may come out by itself. You may become pregnant if the implant comes out by itself. Use a back-up birth control method and call your healthcare professional right away if the implant comes out.
- The need for surgery in the hospital to remove the implant
- Injury to nerves or blood vessels in your arm
- The implant breaks, making removal difficult

• Ectopic Pregnancy

If you become pregnant while using NEXPLANON, you have a slightly higher chance that the pregnancy will be ectopic (occurring outside the womb) than do women who do not use birth control. Unusual vaginal bleeding or lower stomach (abdominal) pain may be a sign of ectopic pregnancy. Ectopic pregnancy is a medical emergency that often requires surgery. Ectopic pregnancies can cause serious internal bleeding, infertility, and even death. Call your healthcare professional right away if you think you are pregnant or have unexplained lower stomach (abdominal) pain.

• Ovarian Cysts

Cysts may develop on the ovaries and usually go away without treatment, but sometimes surgery is needed to remove them.

• Breast Cancer

It is not known whether NEXPLANON use changes a woman's risk for breast cancer. If you have breast cancer now, or have had it in the past, do not use NEXPLANON because some breast cancers are sensitive to hormones.

• Serious Blood Clots

NEXPLANON may increase your chance of serious blood clots, especially if you have other risk factors such as smoking. It is possible to die from a problem caused by a blood clot, such as a heart attack or a stroke.

Serious blood clots can occur within blood vessels of different parts of the body, including the following examples:

- Legs (deep vein thrombosis)
- Lungs (pulmonary embolism)
- Brain (stroke)
- Heart (heart attack)
- Eyes (total or partial blindness)

The risk of serious blood clots is increased in women who smoke. If you smoke and want to use NEXPLANON, you should quit. Your healthcare professional may be able to help.

Tell your healthcare professional at least 4 weeks before if you are going to have surgery or will need to be on bed rest. You have an increased chance of getting blood clots during surgery or bed rest.

- **Other Risks**

A few women who use birth control that contains hormones may get the following conditions:

- High blood pressure
- Gallbladder problems
- Rare cancerous or noncancerous liver tumors

- **Broken or Bent Implant**

Breakage or bending of the implant while it is in your arm may occur due to external forces (such as manipulation of the implant or contact sports). A broken implant may move from the insertion site. If you feel that the implant may have broken or bent while in your arm, contact your healthcare professional.

When should I call my healthcare professional?

Call your healthcare professional right away if you have any of the following conditions:

- Pain in your lower leg that does not go away
- Severe chest pain or heaviness in the chest
- Sudden shortness of breath, sharp chest pain, or coughing blood
- Symptoms of a severe allergic reaction, such as swollen face, tongue or throat; trouble breathing or swallowing
- Sudden severe headache unlike your usual headaches
- Weakness or numbness in your arm, leg, or trouble speaking
- Sudden partial or complete blindness
- Yellowing of your skin or whites of your eyes, especially with fever, tiredness, loss of appetite, dark colored urine, or light-colored bowel movements
- Severe pain, swelling, or tenderness in the lower stomach (abdomen)
- Lump in your breast
- Problems sleeping, lack of energy, tiredness, or you feel very sad
- Heavy menstrual bleeding

What if I become pregnant while using NEXPLANON?

You should see your healthcare professional right away if you think that you may be pregnant. It is important to remove the implant and make sure that the pregnancy is not ectopic (occurring outside the womb). Based on experience with other hormonal contraceptives, NEXPLANON is not likely to cause birth defects.

Can I use NEXPLANON when I am breastfeeding?

If you are breastfeeding your child, you may use NEXPLANON if 4 weeks have passed since you had your baby. A small amount of the hormone contained in NEXPLANON passes into your breast milk. The health of breast-fed children whose mothers were using the implant has been studied up to 3 years of age in a small number of children. No effects on the growth and development of the children were seen. If you are breastfeeding and want to use NEXPLANON, talk with your healthcare professional for more information.

General information about the safe and effective use of NEXPLANON

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. If you would like more information, talk with your healthcare professional. You can ask your healthcare professional for information about NEXPLANON that is written for healthcare professionals.

Manufactured for: Organon USA LLC, a subsidiary of
 **ORGANON & Co.,**
Jersey City, NJ 07302, USA

Manufactured by: N.V. Organon, Oss, The Netherlands, a subsidiary of Organon & Co., Jersey City, NJ 07302, USA

For patent information: www.organon.com/our-solutions/patent/

© 2026 Organon group of companies. All rights reserved.

usppi-og8415-iptx-2601r003

For more information, go to www.nexplanon.com or call 1-844-674-3200.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 01/2026